VOL. 537, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY JANUARY 11, 1991

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

Column Liquid Chromatography

Method for the characterization of stationary phases for the separation of proteins by high-performance liquid chromatography
by R. E. Huisden, J. C. Kraak and H. Poppe (Amsterdam, The Netherlands) (Received August 6th, 1990) 1

Chemical, physical and chromatographic properties of Superdex 75 prep grade and Superdex 200 prep grade gel filtration media
by L. Kågedal, B. Engström, H. Ellegren, A.-K. Lieber, H. Lundström, A. Sköld and M. Schenning (Uppsala, Sweden) (Received August 20th, 1990) 17

Molecular interactions in liquid chromatography
by S. N. Lanin and Yu. S. Nikitin (Moscow, U.S.S.R.) (Received June 9th, 1990) 33

Comparison of retention index scales based on alkyl aryl ketones, alkan-2-ones and 1-nitroalkanes for polar drugs on reversed-phase high-performance liquid chromatography
by R. M. Smith and N. Finn (Loughborough, U.K.) (Received July 31st, 1990) 51

Changes in the enthalpy and entropy of hydroxyl aromatics in reversed-phase liquid chromatography with β-cyclodextrin in the mobile phase
by R. M. Mohseni and R. J. Hurtubise (Laramie, WY, U.S.A.) (Received August 30th, 1990) 61

Comparison of the liquid chromatographic behaviour of selected steroid isomers using different reversed-phase materials and mobile phase compositions
by M. Olsson (Stockholm, Sweden) and L. C. Sander and S. A. Wise (Gaithersburg, MD, U.S.A.) (Received August 27th, 1990) 73

Fatty acid conjugates of chlorinated phenols and their high-performance liquid chromatographic analysis
by B. S. Kaphalia (Galveston, TX, U.S.A.) (Received August 28th, 1990) 85

Separation and identification of free phenolic acids in wines by high-performance liquid chromatography
by G. P. Cartoni, F. Coccioli, L. Pontelli and E. Quattruci (Rome, Italy) (Received July 13th, 1990) 93

High-performance liquid chromatography of red fruit anthocyanins
by J.-P. Goiffon, M. Brun and M.-J. Bourrier (Montpellier, France) (Received August 31st, 1990) 101

Solid-phase reagent containing the 3,5-dinitrophenyl tag for the improved derivatization of chiral and achiral amines, amino alcohols and amino acids in high-performance liquid chromatography with ultraviolet detection
by A. J. Bourque and I. S. Krull (Boston, MA, U.S.A.) (Received August 1st, 1990) . . . 123

Automated phenylthiocarbamyl amino acid analysis of carboxypeptidase/aminopeptidase digests and acid hydrolysates
by R. S. Thoma and D. L. Crimmins (St. Louis, MO, U.S.A.) (Received July 24th, 1990) . 153

(Continued overleaf)

Contents (continued)

Breakthrough curves of insulin, [D-Phe^6]-Gonadotropin-releasing hormone and phenylalanine methyl ester on copolymers of alkylacrylate and divinylbenzene
by P. Slonina and A. Seidel (Berlin, G.D.R.) (Received August 7th, 1990) 167

Alternative mobile phases for reversed-phase high-performance liquid chromatography of peptides and proteins
by B. S. Welinder and H. H. Sørensen (Gentofte, Denmark) (Received July 20th, 1990) . 181

Reversed-phase high-performance liquid chromatographic characterization of acetic acid extracts of the normal and the diabetic human pancreas
by B. S. Welinder and S. Linde (Gentofte, Denmark) (Received July 20th, 1990) 201

Protein–ligand interactions studied on bovine serum albumin with free and polymer-bound Cibacron Blue F3G-A as ligand with reference to affinity partitioning
by G. Johansson and M. Joelsson (Lund, Sweden) (Received August 28th, 1990) 219

High-performance liquid chromatographic analysis of the (cyanoaquo) stereoisomers of several putative vitamin B_{12} precursors
by S. H. Ford, J. Gallery, A. Nichols and M. Shambee (Chicago, IL, U.S.A.) (Received August 30th, 1990) 235

Elimination of peak splitting in the liquid chromatography of the proline-containing drug enalapril maleate
by J. Šalamoun and K. Šlais (Brno, Czechoslovakia) (Received July 30th, 1990) 249

Analysis of ^{55}Fe-labeled hydroxamate siderophores by high-performance liquid chromatography
by R. J. Speirs and G. L. Boyer (Syracuse, NY, U.S.A.) (Received October 2nd, 1990) . . 259

Simultaneous determination of arsenite, arsenate, selenite and selenate in waters using suppressed ion chromatography with ultraviolet absorbance detection
by S. S. Goyal, A. Hafez and D. W. Rains (Davis, CA, U.S.A.) (Received May 30th, 1990) 269

Separation of lanthanides and yttrium as anionic complexes by isocratic ion-interaction chromatography
by E. A. Jones and H. S. Bezuidenhout (Randburg, South Africa) and J. F. van Staden (Pretoria, South Africa) (Received July 24th, 1990) 277

Preseparation of uranium by reversed-phase partition chromatography. Application to the determination of trace elements in nuclear-grade uranium compounds by inductively coupled plasma atomic emission spectrometry
by A. S. Al-Ammar, H. A. Hamid, B. H. Rashid and H. M. Basheer (Baghdad, Iraq) (Received May 15th, 1990) 287

Gas Chromatography

Determination of contributions of different types of solute–sorbent interactions in gas-adsorption chromatography by linear regression of adsorption energies
by O. G. Larionov, V. V. Petrenko and N. P. Platonova (Moscow, U.S.S.R.) (Received July 24th, 1990) 295

Gas chromatographic identification of complex mixtures of halomethanes and haloethanes by using the correlation between their retention and vapour pressure
by T. G. Gerbino and G. Castello (Genova, Italy) (Received July 31st, 1990) 305

Adsorption/thermal desorption for the determination of volatile organic compounds in water
by M. E. Rosen and J. F. Pankow (Beaverton, OR, U.S.A.) (Received May 16th, 1990) . 321

Isomer identification and gas chromatographic retention studies of monomeric cyclic fatty acid methyl esters
by J. A. Rojo and E. G. Perkins (Urbana, IL, U.S.A.) (Received September 7th, 1990) . . 329

Capillary gas chromatographic behavior of stereoisomeric bile acids with a *vicinal* glycol structure by their "mixed" alkylboronate derivatives
by T. Iida and I. Komatsubara (Koriyama, Japan), F. C. Chang (Claremont, CA, U.S.A.) and J. Goto and T. Nambara (Sendai, Japan) (Received August 14th, 1990) 345

Indirect determination of isocyanates by gas chromatography
by C. B. Fanska, T. J. Byerley and J. D. Eick (Kansas City, MO, U.S.A.) (Received July 10th, 1990) 357

Correlation between structure and gas chromatographic behaviour of nitrogen-containing heterocyclic compounds. I. Methyl substitution of pyridopyrimidine derivatives
by O. Papp, Gy. Szász, J. Kökösi and I. Hermecz (Budapest, Hungary) (Received August 21st, 1990) 365

Correlation between structure and gas chromatographic behaviour of nitrogen-containing heterocyclic compounds. II. Alkyl substitution of quinazolone derivatives
by O. Papp, Gy. Szász, L. Őrfi and I. Hermecz (Budapest, Hungary) (Received August 21st, 1990) 371

Correlation between structure and gas chromatographic behaviour of nitrogen-containing heterocyclic compounds. III. Variation of ring size
by O. Papp, Gy. Szász, J. Kökösi and I. Hermecz (Budapest, Hungary) (Received August 21st, 1990) 377

Determination of nine acidic herbicides in water and soil by gas chromatography using an electron-capture detector
by F. Ngan and T. Ikesaki (Sacramento, CA, U.S.A.) (Received August 14th, 1990) . . . 385

Planar Chromatography

Determination of selenium in organic compounds on a silica gel sintered thin-layer chromatographic plate with 2,3-diaminonaphthalene after direct digestion
by R. Hasunuma and T. Ogawa (Kanagawa, Japan), J. Ishii (Tokyo, Japan) and Y. Kawanishi (Kanagawa, Japan) (Received August 7th, 1990) 397

Electrophoresis

Determination of absolute mobilities, p*K* values and separation numbers by capillary zone electrophoresis. Effective mobility as a parameter for screening
by J. L. Beckers, F. M. Everaerts and M. T. Ackermans (Eindhoven, The Netherlands) (Received August 7th, 1990) 407

Isotachophoresis with electroosmotic flow: open *versus* closed systems
by J. L. Beckers, F. M. Everaerts and M. T. Ackermans (Eindhoven, The Netherlands) (Received August 8th, 1990) 429

NOTES

Column Liquid Chromatography

Post-column oligosaccharide dehydrogenase reactor for coulometric detection of malto-oligosaccharides in a liquid chromatographic system
by N. Kiba, K. Shitara and M. Furusawa (Kofu, Japan) and Y. Takata (Katsuta, Japan) (Received August 2nd, 1990) 443

High-performance liquid chromatographic determination of flavonoid glucosides from *Helichrysum italicum*
by P. Pietta, P. Mauri, C. Gardana, R. M. Facino and M. Carini (Milan, Italy) (Received September 3rd, 1990) 449

Isocratic column liquid chromatographic separation of a complex mixture of epicuticular flavonoid aglycones and intracellular flavonol glycosides from *Cistus laurifolius* L.
by T. Vogt and P.-G. Gülz (Köln, F.R.G.) (Received August 7th, 1990) 453

Behaviour of concanavalin A on DEAE-Sephadex and CM-Sephadex
by C. Elias and A. M. Diwan (Pune, India) (Received April 20th, 1990) 460

(Continued overleaf)

Large-scale isolation and purification of human apolipoproteins A-I and A-II
by A. Sigalov, O. Alexandrovich and E. Strizevskaya (Moscow, U.S.S.R.) (Received June 24th, 1990) 464

Preparative high-performance liquid chromatography on chemically modified porous glass. Isolation of acidic saponins from ginseng
by H. Kanazawa, Y. Nagata, Y. Matsushima, M. Tomoda and N. Takai (Tokyo, Japan) (Received August 14th, 1990) 469

High-performance liquid chromatography of substituted trinuclear osmium carbonyl clusters, $Os_3(CO)_{12-n}[P(C_6F_5)_3]_n$ ($n=0, 1, 2$)
by H. G. Ang, W. L. Kwik and W. K. Leong (Singapore, Singapore) (Received August 20th, 1990) 475

Gas Chromatography

Liquid chrystals for the gas chromatographic determination of the stereochemistry of insect sex pheromones
by I. Nesterova, B. Rekhter and G. Roshka (Kishinev, U.S.S.R.) and Z. Witkiewicz (Warsaw, Poland) (Received June 19th, 1990) 482

Simultaneous determination of tributyl phosphate and dibutyl phosphate in spent fuel reprocessing streams by gas chromatography
by Y. Kuno, T. Hina, T. Akiyama and M. Matsui (Ibaraki, Japan) (Received September 11th, 1990) 489

Planar Chromatography

Localization of amino acids on thin-layer chromatograms with acetylacetone–formaldehyde reagent
by M. B. Devani, C. J. Shishoo, S. A. Shah, K. P. Soni and R. S. Shah (Ahmedabad, India) (Received August 31st, 1990) 494

LETTERS TO THE EDITOR

Pitfalls in the choice of isotherms for the calculation of band profiles in preparative chromatography
by M. Czok and G. Guiochon (Knoxville and Oak Ridge, TN, U.S.A.) (Received October 17th, 1990) 497

Pitfalls in the choice of isotherms for the calculation of band profiles in preparative chromatography. A reply
by L. R. Snyder (Lafayette, CA, U.S.A.) and G. B. Cox (Indianapolis, IN, U.S.A.) (Received September 25th, 1990) 507

Detection of choline and acetylcholine by high-performance liquid chromatography. Limitations, pitfalls, sample preparation
by E. Haen (Munich, F.R.G.), H. Hagenmaier (Tübingen, F.R.G.) and J. Remien (Munich, F.R.G.) (Received August 27th, 1990) 514

Author Index 520

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)